

TABLE A1.1. Proxima Cen Log of Observations

mJD	Exposure Time	Average Intensity	mJD	Exposure Time	Average Intensity
8704.858	93.60	3972.96	8858.714	92.40	4038.52
8704.871	66.45	3988.01	8858.733	64.35	4049.20
8713.824	92.55	3971.07	8866.820	92.40	4021.89
8713.838	66.45	4001.50	8866.837	79.35	4019.69
8723.807	92.40	4015.08	8876.226	92.40	3995.54
8723.820	66.45	4016.87	8886.134	92.25	4027.40
8736.396	93.45	4044.73	8886.151	79.35	4027.74
8736.410	67.50	4055.95	8896.979	92.40	4051.47
8751.390	92.40	3986.84	8896.996	79.50	4057.43
8751.404	66.45	3995.15	8906.083	92.55	4192.68
8756.509	92.40	3999.22	8906.101	79.35	4045.50
8756.524	66.30	4007.55	8916.583	92.40	3911.94
8759.860	92.40	4026.46	8916.597	79.50	3912.24
8759.873	66.45	4016.65	8977.539	92.40	4095.24
8769.707	92.40	4044.77	8988.275	92.25	3966.21
8769.720	66.45	4036.54	8988.293	79.35	3960.82
8783.702	92.40	3981.93	8999.116	92.25	3870.63
8783.715	66.60	3997.98	8999.134	79.50	3874.66
8790.130	92.40	3992.80	9008.892	92.25	3873.44
8790.143	67.35	4002.20	9008.909	79.35	3886.46
8813.522	92.25	4072.92	9091.916	92.25	3849.56
8813.541	62.40	4077.23	9091.933	79.35	3844.65
8813.662	92.40	4075.28	9104.975	92.40	3938.08
8813.681	64.50	4081.69	9104.992	79.20	3944.91
8813.723	92.40	4080.03	9115.683	93.30	4042.42
8813.742	64.50	4081.36	9115.698	79.35	4053.37
8819.420	92.25	4064.21	9126.589	91.35	4098.67
8819.438	79.35	4057.65	9126.603	79.35	4112.53
8823.371	92.25	4075.47	9136.355	91.35	4120.11
8823.390	64.50	4064.61	9136.369	79.35	4134.43
8823.432	92.40	4030.04	9154.721	82.35	3963.77
8823.451	64.50	4032.01	9154.736	73.35	3963.73
8827.522	92.40	4010.21	9163.753	83.25	3856.28
8827.539	79.35	3993.68	9163.768	73.20	3909.88
8830.469	92.25	3973.40	9210.749	84.30	4136.98
8830.479	64.50	3974.33	9210.767	77.25	4137.28
8830.529	92.25	3967.63	9227.202	161.25	4133.59
8830.548	64.50	4013.47	9227.214	106.20	4129.79
8845.197	92.40	3993.66	9227.225	106.20	4160.07
8845.216	64.50	4000.27	9237.104	131.25	3989.11
8845.259	93.45	4023.71	9237.116	106.20	3990.16
8845.278	64.50	4013.82	9237.127	104.40	3989.45
8845.330	92.40	4000.19	9265.502	131.25	3922.78
8845.347	79.50	4003.18	9265.514	106.65	3965.31
8855.975	92.40	4008.64	9265.525	101.25	4032.79
8855.992	79.35	4004.59	9367.956	131.25	4330.42
8857.844	92.40	4015.44	9367.968	104.25	4166.76
8857.862	64.50	4010.45	9367.979	103.20	4102.59

TABLE A1.1. (continued)

mJD	Exposure Time	Average Intensity	mJD	Exposure Time	Average Intensity
9439.099	160.50	4029.86	10097.187	712.05	4056.11
9439.111	104.25	4032.52	10097.206	657.75	4057.65
9439.122	104.25	4035.69	10118.897	103.95	4146.12
9448.744	130.35	4012.61	10118.907	106.95	4153.15
9448.756	103.35	4015.98	10118.911	110.85	4154.69
9448.767	103.20	4017.74	10147.225	105.00	4054.12
9458.461	130.20	4018.50	10147.235	106.95	4058.27
9458.473	103.20	4018.41	10147.239	109.95	4097.83
9458.484	103.35	4017.34	10208.929	684.00	4186.41
9467.510	131.25	4090.33	10208.944	714.90	4178.43
9467.522	104.25	4083.18	10208.959	657.90	4174.19
9467.529	109.20	4080.91	10263.709	685.05	4038.82
9476.629	131.25	4073.87	10263.726	713.85	4039.39
9476.641	103.20	4083.48	10263.745	658.05	4038.92
9476.652	97.65	3651.69	10335.487	174.90	3929.83
9924.038	686.10	3994.89	10335.498	174.90	3929.85
9924.055	714.00	4005.25	10335.509	175.05	3935.00
9924.074	657.90	4005.32	10335.514	177.90	3936.73
9924.102	107.85	3990.95	10372.576	172.95	4095.93
9924.112	106.95	3995.77	10372.587	173.85	4104.08
9924.123	106.95	3992.63	10372.599	170.85	4105.24
9924.126	109.95	3996.53	10372.603	177.90	4112.94
9951.202	108.00	4006.80	10556.684	173.85	4077.49
9951.212	106.80	4010.80	10556.684	173.85	4077.49
9951.223	108.90	4032.56	10556.696	170.70	4098.21
9951.226	109.95	4050.83	10556.696	170.70	4098.21
9985.447	684.00	4010.56	10556.707	172.80	4083.36
9985.464	712.20	4003.36	10556.707	172.80	4083.36
9985.484	657.90	4023.77	10556.711	177.90	4089.84
10000.199	106.95	3971.31	10708.050	172.90	4033.45
10000.209	107.10	3968.96	10708.059	176.90	4044.00
10000.220	102.90	3970.80	10708.070	174.70	4039.26
10000.224	109.95	3974.83	10708.075	177.90	4039.48
10082.549	106.95	3907.21	10736.169	172.90	3919.48
10082.559	106.05	3910.19	10736.180	174.00	3936.21
10082.570	103.95	3908.66	10736.191	171.90	3913.60
10082.573	110.10	3916.68	10736.196	177.90	3912.12
10097.170	684.90	4043.80			

TABLE A1.2. Barnard's Star Log of Observations

mJD	Exposure Time	Average Intensity	mJD	Exposure Time	Average Intensity
9025.634	80.4	15065	9439.92	39.45	15054.7
9025.649	24.3	15046.7	9439.927	35.25	15053.4
9025.658	24.3	15064.8	9473.56	90.3	14989.1
9088.103	50.25	15106.1	9473.575	38.4	15012.9
9088.117	24.3	15116.5	9473.581	35.4	15022.7
9088.127	24.45	15115.3	9487.853	37.35	15007.9
9102.763	50.25	15069.3	9487.86	34.35	15012.2
9102.777	24.3	15091.7	9506.676	91.2	14955.1
9102.788	24.15	15093.1	9506.691	36.3	14945.1
9118.631	51.15	14978.7	9506.698	33.3	14947
9118.645	24.3	14994.9	9522.698	93.15	14870.4
9118.658	24.3	14979.9	9522.713	33.15	14880.3
9128.672	49.35	14895.8	9522.72	34.05	14886.2
9128.686	24.45	14914.1	9536.508	92.1	14918.5
9128.699	24.45	14915.4	9536.523	34.05	14901.1
9131.751	49.35	14880.2	9536.53	33.9	14931.1
9131.765	24.45	14895.7	9566.407	122.1	15001.8
9131.777	24.3	14905.7	9566.422	34.05	14967
9184.57	122.25	15045.9	9566.428	33	14977.1
9184.585	33.3	15028.5	9581.286	122.1	15033
9184.591	32.25	15034.9	9581.301	34.05	15016.3
9216.568	91.35	15047.8	9581.306	34.05	15033.6
9216.583	34.35	15021.6	9594.493	121.95	15007.1
9216.586	33.3	15035.4	9594.508	37.05	15028.9
9425.762	120.3	15082	9594.515	35.1	15050.2
9425.777	35.25	15070.2	9609.505	121.8	14962.2
9425.781	35.25	15080.4	9609.52	36	14965.5
9439.905	91.2	15076.4	9609.527	35.85	14982.9
9622.11	121.95	14952.7	9962.146	76.95	14996.1
9622.126	36	14925.8	9962.157	75	15001.7
9622.132	35.85	14957.6	9994.169	117	14977.5
9636.119	122.1	14943.7	9994.181	76.95	15020.3
9636.134	37.05	14916.4	9994.192	75	15035.7
9636.141	36	14928.4	10024.262	118.05	14974.8
9653.682	121.95	14918.7	10024.274	78	14997.2
9653.697	37.95	14923.1	10024.285	75	15002.1
9653.703	34.95	14938	10117.955	13.2	14986
9663.199	122.1	14946.8	10117.968	75.9	14986.7
9663.214	37.95	14967.6	10117.979	48.48	14981.9
9663.221	34.05	14984.8	10143.754	77.1	14982.8
9757.843	123	15002.6	10143.768	75.75	14978
9757.858	36.15	14984.8	10143.778	75	14980.8
9757.864	33.9	14997.5	10167.077	115.95	14968.8
9777.147	92.1	14937.5	10167.077	115.95	14968.8
9777.162	36	14910.5	10167.077	115.95	14968.8
9777.169	34.95	14929.3	10167.1	73.95	14985.5
9916.557	116.85	14927.3	10198.508	116.85	14903.2
9916.57	76.95	14925.7	10198.52	76.95	14921.4

TABLE A1.2. (continued)

mJD	Exposure Time	Average Intensity	mJD	Exposure Time	Average Intensity
9916.581	75	14919.7	10198.531	75.9	14928.4
9935.324	117	14882.6	10237.311	118.05	14914
9935.336	77.85	14917	10237.324	80.85	14942.5
9935.347	76.05	14934.9	10237.335	56.73	14931
9962.134	117	14984.8			

**Photometry of Proxima Centauri and Barnard's Star
Using HST Fine Guidance Sensor 3:
A Search for Periodic Variations¹**

G. Fritz Benedict², Barbara McArthur², E. Nelan³, D. Story⁹, A. L. Whipple¹⁰, P. J. Shelus², W.
H. Jefferys⁴, P.D. Hemenway⁵, Otto G. Franz⁶, L. H. Wasserman⁶, R. L.
Duncombe¹¹, Wm. van Altena⁷, and L.W. Fredrick⁸

ABSTRACT

We have observed Proxima Centauri and Barnard's Star with *Hubble Space Telescope* Fine Guidance Sensor 3. Proxima Centauri exhibits small-amplitude, periodic photometric variations. Once several sources of systematic photometric error are corrected, we obtain 2 milli-magnitude internal photometric precision. We identify two distinct behavior modes over the past four years: higher amplitude, longer period; smaller amplitude, shorter period. Within the errors one period ($P \sim 83^d$) is twice the other. Barnard's Star shows very weak evidence for periodicity on a timescale of approximately 130 days. If we interpret these periodic phenomena as rotational modulation of star spots, we identify three discrete spots on Proxima Cen and possibly one spot on Barnard's Star. We find that the disturbances change significantly on time scales as short as one rotation period.

¹Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555

²McDonald Observatory, University of Texas

³Space Telescope Science Institute

⁴Astronomy Dept., University of Texas

⁵University of Rhode Island

⁶Lowell Observatory

⁷Astronomy Dept., Yale University

⁸Astronomy Dept., University of Virginia

⁹McDonald Observatory, University of Texas. Now at Jackson & Tull.

¹⁰McDonald Observatory, University of Texas. Now at Allied-Signal Aerospace.

¹¹Aerospace Engineering, University of Texas

Subject headings: stars: flare — stars: individual (Proxima Centauri, Barnard’s Star)
— stars: late-type — stars: rotation — stars: spots

1. Introduction

We present photometry of Proxima Cen and Barnard’s Star, results ancillary to our astrometric searches for planetary-mass companions (Benedict et al., 1997a). Our observations were obtained with Fine Guidance Sensor 3 (FGS 3), a two-axis, white-light interferometer aboard the *Hubble Space Telescope (HST)*. Bradley et al., 1991 provide an overview of the FGS 3 instrument and Benedict et al., 1994a describe the astrometric capabilities of FGS 3 and typical data acquisition strategies. Benedict et al., 1993 assessed FGS 3 photometric qualities and presented the first evidence for periodic variability of Proxima Cen. This latter result was based on 212 days of monitoring. Subsequent data exhibited a period of variation very nearly twice the original (Benedict et al., 1994b). Since that report, we have obtained 14 additional data sets for Proxima Cen and 12 new sets for Barnard’s Star. The primary value of these observations lies in their precision, not in their temporal span or aggregate numbers. We have previously determined that a 90 sec observation obtained with FGS 3 has a $1 - \sigma$ precision of 0.001 magnitude at $V = 11$ (Benedict et al., 1993), in the absence of systematic errors.

In this paper we discuss the data sets and assess systematic errors, including background contamination and FGS position-dependent photometric response. We also present a revised photometric flat field. We then exhibit and analyze light curves for Proxima Cen and Barnard’s Star. We find weak evidence for periodic variations in the brightness of Barnard’s Star. However, Proxima Cen exhibits significant periodic photometric variations, with changes in amplitude and/or period. We next interpret these variations as rotational modulation of chromospheric structure (star spots and/or plages), and conclude with a brief comparison to other determinations of the rotation rate of Proxima Cen. Tables 1 and 2 provide aliases and physical parameters for our two science targets.

We use the term ‘pickle’ to describe the total field of view of the FGS. The instantaneous field of view of FGS 3 is a 5×5 arcsec square aperture. Figure 1 shows a finding chart for the Barnard’s Star reference frame in the FGS 3 pickle as observed on 6 August 1994. Benedict et al., 1993 contains a finding chart for the Proxima Cen reference frame.

2. Data Reduction

2.1. The Data

All position and brightness measurements from FGS 3 are comprised of series of 0.025 sec samples (e.g., 40Hz data rate), of between 20 and 120 sec or ~ 600 sec duration. Each FGS

contains four photomultipliers, two for each axis. We sum the output of all four to produce our measurement, S , the average count per 0.025 sec sample, obtained during the entire exposure. The coverage for both targets suffers from extended gaps, due to *HST* pointing constraints (described in Benedict et al., 1993) and other scheduling difficulties. The filter (F583W) has a bandpass centered on 583 nm, with 234 nm FWHM.

For Proxima Cen the data now include 152 shorter exposures secured over 4 years (March 1992 to October 1997) and 15 longer exposures (July 1995 to July 1996). Each orbit contains from two to four exposures. The longest exposure times pertain only to Proxima Cen observations obtained within Continuous Viewing Zone (CVZ) orbits. These specially scheduled orbits permit ~ 90 minutes on field, during which Proxima Cen was not occulted by the Earth. Appendix 1.1 gives times of observation, exposure times, and average counts, S , for all Proxima Cen photometry.

Barnard’s Star was monitored for three years (February 1993 to April 1996), and observed three times during each of 35 orbits. Exposures range between 24 and 123 seconds duration. Appendix 1.2 gives times of observation, exposure times, and average counts, S , for all Barnard’s Star photometry.

2.2. Background Light

We first noted that background contamination might be an issue while assessing the use of astrometric reference stars for photometric flat-fielding. These stars are typically far fainter than the primary science targets. Using them to flat field the Proxima Cen and Barnard’s Star photometry introduced a strong one-year periodicity (and considerable noise, since they are fainter stars). This problem was not identified in Benedict et al., 1993, since we had access to data spanning less than two-thirds of a year. Figure 2 shows S for two faint reference stars in the Barnard field plotted against angular distance from the Sun. These stars appear brightest when closest to the Sun. Zodiacal light is a source whose brightness depends on the sun-target separation. The fitting function in Figure 2 is

$$I = A + B \sin\left(\frac{\theta}{2}\right), \quad (1)$$

chosen to produce a minimum contribution at $\theta = 180^\circ$. We find $A = 137.1 \pm 0.4$ and $B = -4.1 \pm 0.5$ counts per 25ms for an average exposure time of 100 sec. At a 60° elongation the contamination amounts to $V = 22.5 \pm 0.3$ arcsec $^{-2}$. The Barnard field is at ecliptic latitude $\beta = +27^\circ$. From a tabulation in Allen (1972) we calculate a signal equivalent to $V = 22.1$ arcsec $^{-2}$ for zodiacal light at 60° elongation and ecliptic latitude $\beta = +30^\circ$. The agreement supports our identification of this background source.

We present in Figure 3 an average S for these two Barnard reference stars plotted as a function of time, uncorrected and corrected for background. These data have been flat fielded using the time-dependent response function discussed in section 2.3.2 (equation 2). Note the

reduction in the amplitude of the scatter for the corrected photometry. Presuming Zodiacal Light as the source, contamination levels are even less for the Proxima Cen observations at ecliptic latitude $\beta = -44^\circ$, introducing a maximum systematic error of 0.0007 magnitude for a 100 sec observation. We conclude that the effects of this component of the background are insignificant for Proxima Cen and Barnard’s Star photometry.

Should background determination become more important in the future, we note that during an intra-orbit observation sequence the PMT are never turned off. Hence, the *HST* data archive contains PMT measurements taken during slews from one star to the next. The astrometric reduction pipeline at the Space Telescope Science Institute has been modified to provide these background data automatically.

2.3. Photometric Flat Fielding

We explore two kinds of flat fielding; position- and time-dependent. We first assess whether or not flat-field corrections are necessary, and, if so, determine their functional form.

2.3.1. Position-dependent Photometric Response

Having discovered that background variations contaminate the photometry of faint astrometric reference stars, we required an alternative source for flat field data. To maintain the astrometric calibration of FGS 3, a star field in M35 has been measured roughly once per month for the last four years. Whipple et al., 1995 describe this continuing astrometric Long-term Stability (LTSTAB) test. The field, on the ecliptic, and, hence, always observed in one of two orientations (Fall or Spring) flipped by 180° , contains bright stars for which background contamination is negligible. However, an initial application of a time-dependent flat field based on bright M35 stars also introduced a strong one-year periodicity.

The positions of the three M35 LTSTAB stars within FGS 3 are shown in Figure 4 . The ‘eye’ is bordered by the pickle edge at the two nominal rolls for this field. The central circle (diameter $\sim 3'8$) is accessible by the FGS 3 instantaneous aperture for any HST roll.

Figure 5 (bottom) presents normalized intensities ($I = S(t)/S_{av}$, where S_{av} is determined from the entire run of data) for the three LTSTAB stars as a function of time. The variation of each star has first been modeled by a linear trend. The parameters, intercept (I_o) and slope (I'), are given in Table 3. The resulting residuals (Figure 5, top) have been modeled with a sine wave. while constraining $P = 365^{\text{d}}.25$ days. The residuals have a square-wave periodic structure because, rather than a range of spacecraft rolls, there are only two orientations. The resulting parameters and errors are given in Table 3. In Figure 7 we plot the amplitude of this side-to-side variation against radial distance from the pickle center. For the M35 stars. the further the star

from the pickle center, the larger the roll-induced variation. Figure 7 includes several other one year period amplitudes; a preliminary result for GJ748 ($V \sim 11.1$, ecliptic lat $\beta \sim +23^\circ$) always observed in the center of the pickle, the Barnard reference star photometry from Figure 3 corrected for background, and photometry of the brightest reference star in the Barnard field (Figure 1, star 36, $V \sim 11.5$). Figure 7 suggests that within the inscribed circle of Figure 1 ($r < 180''$), position-dependent photometric response variations should be less than 0.002 magnitude.

We have also identified one high spatial frequency position-dependent flat field component for FGS 3. Light curves for two of the Barnard reference stars evidenced sudden decreases in brightness with subsequent return to previous levels. The decrease for reference star 34 was 29%; for star 36, 17%. Shown in Figure 8, both decreases occurred in the same location within FGS 3, very near the -Y edge. The pickle X, Y of the center of this area is $(X, Y) = -25, 627$. We estimate the size of the low-sensitivity region to be $\sim 10 \times 10$ arcsec. Additionally, Proxima Cen reference star observations acquired one year prior to the Barnard’s Star reference star observations and within a few arcsec of this position showed no decrease, providing additional evidence that FGS 3 is not suitable for wide-field, precise faint star photometry.

Evidence that the photometric response may vary locally and randomly with time dissuades us from mapping a position-dependent flat field over the entire pickle. However, for bright stars ($V < 11$) observed within ~ 20 arcsec of the pickle center (Figure 1), these identified systematics should produce very little effect. *All Proxima Cen and Barnard’s Star observations were secured within 15 arcsec of the pickle center.*

2.3.2. Time-dependent Photometric Response

Figure 5 indicates that FGS 3 has become less sensitive with age. For all three LTSTAB stars the linear trends (I' , Table 3) agree within the errors. The apparent 1% drop in sensitivity over 1000 days requires confirmation. Figure 6 presents the time varying normalized intensity for two other astrometric program stars observed with FGS 3, GJ 623 and GJ 748. Both M dwarfs were observed in pickle center. Comparing the I' in Table 3 and Table 4, the rate of decline in brightness for GJ 623 and GJ 748 is identical (within the errors) to that seen in the M35 stars.

A final concern is that the rate of decline of PMT sensitivity might vary with wavelength. The M35 stars (stars 547, 500, and 312 in the catalog of Cudworth, 1971) have $0.12 < B - V < 0.49$, while GJ 623 and GJ 748 have $B - V \simeq +1.5$. There appears to be no dependence on color.

The weighted average for five stars from three different fields yields

$$FF = 1.131 \pm 0.006 + (1.30 \pm 0.06) \times 10^{-5} mJD \quad (2)$$

as the temporal photometric flat field for the pickle center.

As an additional test of the reality of this sensitivity decrease, we note that the intensity data

for the two astrometric reference stars in the Barnard’s Star field shown in Figure 3 have been flat-fielded with equation 2. Thus, a total of seven stars from four different fields show similar brightness trends, adequate evidence for a sensitivity loss in FGS 3.

2.4. Photometric Calibration

All magnitudes presented in this paper are provisional, since a final calibration from F585W to V is not yet available. If magnitudes are given, they are derived through

$$V = -2.5 \log(S) + 20.0349 \quad (3)$$

with no color term, where S is the average count per 0.025 sec sample, summed over all four PMT. No results are based on these provisional calibrated magnitudes. They are provided only as a convenience.

2.5. Summary: Photometric Error and Photometric Precision

We have identified sky background (Zodiacal Light), within-pickle response variations, and time-dependent sensitivity variations as contributing sources of systematic error for our photometry. Since our science targets, Proxima Cen and Barnard’s Star, are bright, the effect of Zodiacal Light is at most 0.001 magnitude. Since we observe these stars only in the pickle center, spatially-induced variations are reduced to about 0.001 magnitude, our claimed per-observation precision at $V \sim 11$. A weighted average of the temporal response of five stars in three fields provides a very precise flat field whose slope error could introduce at most 0.001 magnitude systematic error over 1000 days. (Since we are doing only differential photometry we ignore the zero point error in the flat field.) Combining these sources of error yields a per-observation precision of 0.002 magnitude.

3. Photometric Results

We apply the flat field (equation 2) to the Appendix 1.1 and Appendix 1.2 S values and plot (Proxima Cen, Figure 9; Barnard’s Star, Figure 10) the resulting intensities as function of modified Julian Date (JD - 2440000). Our coverage in time is not uniform for either target. There are extended gaps in coverage, some due to the *HST* solar constraint (no observations permitted closer than $\pm 50^\circ$ to the Sun). The largest gap, in 1994 for Proxima Cen, was due to an awkward transition from Guaranteed Time Observations to Guest Observer status and a hiatus due to suspected equipment problems.

3.1. Trends in Brightness

For Proxima Cen the solid line in Figure 9 indicates an overall trend of increasing brightness with time. In units of normalized intensity the rate of change of brightness ($1.63 \pm 0.37 \times 10^{-5}$) is similar to that of the adopted flat field (equation 2). For Barnard’s Star (Figure 10) the slope of the upward trend in units of normalized intensity is $+0.91 \pm 0.18 \times 10^{-5}$, again, suspiciously similar in absolute value to the adopted flat-field relation (equation 2). Since seven stars from four different fields exhibit the sensitivity decrease described by the flat field, the Proxima and Barnard upward trends are unlikely to be a flat-field artifact.

A final caveat: Proxima Cen and Barnard’s Star are somewhat redder (Tables 1 and 2) than GJ 623 and GJ 748. If the sensitivity loss varies with wavelength (e.g., more sensitivity loss for blue than for red wavelengths), it would have to be a very steeply dependent function, showing no effect at $B - V = +1.5$.

3.2. Proxima Cen

The flat-fielded photometry for each exposure in each orbit appears in Figure 9. The period and amplitude variations evident in Figure 9 will be discussed in section 4. Our total time on target, obtained by summing the exposure times in Appendix A.1, was 6^h6. Proxima Cen is a flare star (V654 Cen) and these data contain exposures ‘contaminated’ by stellar flares (marked F1 - F4 in Figure 9). We identified these events by inspecting the 40Hz photometric data stream for each observation. An example of flare contamination (including a light curve) can be found in Benedict et al., 1993, which discusses a slow, relatively faint ($\Delta V < -0.10$), and multi-peaked flare on mJD 8906 (F1 in Figure 9). An explosive flare on mJD 9266 ($\Delta V \sim 0^{\text{m}}6$ in one second, F3 in Figure 9) produced astrometric changes in Proxima Cen, analyzed in detail by Benedict et al., 1997b. This spectacular event provided the motivation for the subsequent CVZ observations (cz in Figure 9), each permitting 30 minutes of monitoring for flares. The F4 event at mJD 9368 had a relatively small amplitude ($\Delta V \sim -0.13$), but lasted throughout the entire 130^s exposure, hence its large effect on the exposure. Walker (1981) predicts a flare with intensity similar to F3 once every 31 hours. Thus, while disappointing, it is not surprising that we captured none as bright as the F3 event in our additional 2.5 hours of CVZ on-target monitoring. It may be significant that we saw any flares at all, since even the small amplitude events have only a 60% chance of occurring during our total monitoring duration. We will discuss this further in Section 5.1.

Individual observations secured within an orbit and not affected by flaring exhibit an internal consistency at the 0.002 magnitude level.

3.3. Barnard’s Star

The flat-fielded photometry for each of the three Barnard’s Star exposures acquired within each orbit appears in Figure 10. Note that the time scale is exactly that used for Figure 9 to facilitate comparison. Again, note that those observations secured within an orbit exhibit an internal consistency at the 0.002 magnitude level. We find variations within each orbit, but no obvious flaring activity in the Barnard’s Star results. Possible period and amplitude variations in the Barnard’s Star data will be discussed in section 4.2.2.

The scatter within each orbit in Figure 10 is somewhat larger than the previously determined (Benedict et al., 1993) 0.001 magnitude measurement precision. In particular we inspected the observations on mJD 9935 and 9994 and found only a slight upward slope during the first observation on each date. Since the majority of first observations within each orbit are lower, this intra-orbit scatter is most likely an instrumental effect, amounting to about 0.001 magnitude. The first observation low bias is sometimes seen in the Proxima Cen data (Figure 9). Leaving all first observations uncorrected will only slightly enlarge the formal errors for our per-orbit means.

4. Analysis

For subsequent analyses of Proxima Cen, we removed the flare contributions by subjecting the per-orbit average to a pruning process. All exposures obtained during each orbit are presented in Figure 9. If one exposure differs by more than $2.5 - \sigma$ from the mean for that orbit, it is removed and the mean recalculated. This process results in 71 normal points with associated dispersions (the standard deviations calculated for 2, 3, or 4 exposures in each orbit) for Proxima Cen. No exposures were removed from the Barnard’s Star series, since no intra-orbit points (shown in Figure 10) violated the $2.5 - \sigma$ criterion. The resulting per-orbit average S values are presented as direct light curves in Figure 13 (Proxima Cen) and Figure 16 (Barnard’s Star). Forming these normal points provides per-orbit photometric precision better than 0.002 magnitude for the following analyses.

4.1. Lomb-Scargle Periodograms

From Figures 9 and 10 we suspect that there are periodic variations in both the Proxima Cen and Barnard photometry. To obtain a preliminary identification of these periodicities we produce Lomb-Scargle Periodograms (Press et al., 1992) from the per-orbit normal points presented in Figure 13 (Proxima Cen) and Figure 16 (Barnard’s Star).

The most statistically significant period in the Proxima Cen periodogram (Figure 11) is at $P \sim 83^d$, with a false-positive probability less than 0.1%. The very small peak at $P \sim 42^d$ indicates the relative strengths of the period derived from the first 212 days (Benedict et al.,

1993) relative to the higher amplitude $P \sim 83^d$ period. The false-positive probability for the period derived in that paper from only the first 212 days was $\sim 1\%$. Since the periodogram provides no results for very short periods, we have some concern that we are undersampling a more rapid variation. We can rule out a range of periods $2^d < P < 20^d$ from detailed inspection of clusters of data near mJD 8840 (Figure 13), where we had a series of closely-spaced (in time) observations (see Appendix 1.1).

The periodogram for Barnard’s Star is shown in Figure 12. The strongest peak (at $P \sim 130^d$) has a 10% false-positive probability. We have much less compelling evidence of variability for Barnard’s Star than for Proxima Cen.

4.2. Light Curves

4.2.1. Proxima Cen Light Curve

Given strong support for a periodic variation (periodogram, Figure 11) and for an overall trend in the brightness (Figure 9), we model the per-orbit average variations seen in the direct light curve (Figure 13) with a sin function and trend

$$I = I_o + I't + A \sin\left(\left(\frac{2\pi}{P}\right)t + \phi\right), \quad (4)$$

To reconcile the earlier results (Benedict et al., 1993) with the newer data, we first attempted to model the entire light curve with only two distinct segments, grouping segments B, C, and D together. From the earliest data (segment A) the Proxima Cen photometric variations are characterized by a shorter period and smaller amplitude. Later data are best fit with a longer period and larger amplitude variation, as evidenced by the periodogram (Figure 11). Parameters for these fits are listed A and BCD in Table 4 (lines 1 and 2). We find $P_{BCD}/P_A = 1.97 \pm 0.04$.

Noting very large residuals for segment C, we next explored the possibility that Proxima Cen repeats a low to high amplitude cycle by fitting the four segments (A - D, Figure 13) with the same model (equation 4). The parameters for these fits are presented in Table 4 (lines 1 and 3 - 5). Within the errors, $P_A = P_C$ and $P_B = P_D$, with $P_D/P_C = 1.99 \pm 0.02$. A reduction in the number of degrees of freedom by 17% (fitting 71 data points with twenty, rather than ten parameters), reduced the residuals by $\sim 30\%$. This relative improvement is some support for alternating high and low amplitude states. It is also evident that segments A and C have very nearly half the period of segments B and D.

Figure 14 contains phased light curves for the four segments labeled in Figure 13. In the top panel we show that the phase shift required to align the two long-period segments is small ($\Delta\phi = +0.11$) for the period $P = 83.5^d$ suggested by the periodogram (Figure 11). We have shifted segment D down by $\Delta S = -104.4$ counts. The bottom panel of Figure 14 shows a phased

light curve for the two shorter period segments (A and C), phased also to $P = 83.5^d$. Shifts in phase and intensity to achieve alignment are indicated in the figure. The clean double sin wave also demonstrates that the low-amplitude segments, A and C, have half the period of the higher-amplitude, longer-period segments, B and D.

Finally, Figure 15 is used to demonstrate that the same low-amplitude, short-period variations seen in segments A and C may also be present in segments B and D. We fit a sin wave to the phased B and D light curve in the lower panel of Figure 15, constraining the period to one cycle. The top panel of Figure 15 shows the residuals to that fit. These residuals are then fit with a sin wave with the period constrained to one-half cycle. Comparing with the bottom panel of Figure 14, we find a similar double sin wave, nearly identical phase, and an amplitude ($A = 25 \pm 8$) close to that reported for segments A and C in Table 5.

4.2.2. *Barnard’s Star Light Curve*

We turn now to the per-orbit average photometry of Barnard’s Star. Figure 16 (bottom) contains the per-orbit average direct light curve. The error bars are $1 - \sigma$, obtained from the dispersion of the three observations within each orbit on each date (Figure 10). Residuals to a linear trend are presented in the top panel of Figure 16. The sin wave fit to these residuals was constrained to have the most significant period detected in the Figure 12 periodogram, $P = 130^d.4$. Figure 17 contains a light curve for the trend-corrected Barnard’s Star photometry of Figure 16 (top), phased to $P = 130^d.4$. The phased light curve is far less clean than for Proxima Cen. The periodogram and Figures 16 and 17 provide only weak evidence for periodic variation, primarily due to the poor sampling.

5. Discussion of Photometric Results

Instruments can impress spurious periodicities on data (Kristian, 1991). It is comforting that we find for all segments of either data set that $P_{Barn} \neq P_{Prox}$.

Stars have local imperfections in their atmospheres (e.g., the Sun, Zirin 1988). Stars other than the Sun have been shown to be spotted, photometrically (dwarf M stars; Kron, 1952) and spectroscopically (e.g., Hatzes, 1993; Neff et al., 1995). Other M stars have been shown to have spots, both dark (Bouvier et al., 1995) and bright (α Ori, Gilliland & Dupree, 1996).

A spot on a rotating star is a model rich in degrees of freedom. Spots can be bright (plages) or dark (see Pettersen et al., 1992 for a discussion of the choice between dark spots on a bright background or bright spots on a dark background). Spots can wax and wane in size, driving the mean brightness level of a star up or down (Vrba et al., 1988). Spots can migrate in latitude, which, when coupled with presumed differential rotation, can change the phasing and perceived

rotation period. Spots are thought to migrate up or down (relative to the star center) within the magnetosphere (Cameron & Campbell, 1993), inducing perceived period changes. In the following sections we shall interpret the variations seen in Figures 13 and 16 as rotational modulation of spots or plages.

5.1. Spots on Proxima Cen

If we assume a fundamental rotation period $P = 83.5^d$, then variations in the amplitude (Figure 13) could be due to spot/plage changes. With the sparse set of single-color photometric data presented in Figure 13 we have made no effort to quantitatively model spots (c.f. Neff et al., 1996). The period and amplitude changes can be qualitatively modeled using plages and spots, but require the disappearance of a feature or a major change in feature size or temperature in less than one rotation period (e.g., the A to B segment transition seen in Figure 13).

Segments B and D (Figure 14, top) require a single large or darker spot that reduces the average brightness of Proxima Cen by $\Delta V \sim 0.03$. This feature was not present in segment A and disappeared during segment C. To phase segments B and D we applied a shift of $\Delta\phi = 0.1$ radians. Thus, the spot site lagged behind the fundamental rotation period by about 5° over the end of B to start of D time separation of $\sim 700^d$. Whether due to latitude migration coupled with differential rotation or from changes of height within the magnetosphere is unknown. If due to differential rotation, then either the spot moved very little in latitude or differential rotation on Proxima Cen is several orders of magnitude less than the Sun (Zirin 1988).

Segments A and C exhibit smaller amplitude variations with a period almost exactly half that found for segments B and D. These segments (see the phased light curves in Figure 14, bottom and Figure 15, top) could be produced by two smaller spots spaced 180° in longitude, carried around by the fundamental 83.5^d rotation period, and persisting through all segments A to D. These two spots produce a $\Delta V \sim 0.01$. One of these small or less dark (warmer) spots lies at nearly the same longitude as the prominent spot seen in segments B and D. The other lies near the center of the brighter hemisphere in segments B and D.

From Figure 13 we note that the peaks in segments B and D were brighter. In segment B the minima were deeper, implying a darker (cooler) spot. To increase the amplitude of the maxima in segments B and D requires the existence of plages, or that the hemisphere not containing the single large spot became brighter due to spot changes. One of the pair of spots, associated with the brighter hemisphere ($\phi \sim 0.3$, Figure 15, top), does have a shallower minimum than the other of the pair at $\phi = 0.8$. If these are the same spots responsible for the variation in segments A and C, then the spot at $\phi \sim 0.3$ did have a shallower minimum in segments B and D (compare Figure 14, bottom and Figure 15, top). However, that spot did not become less dark by enough to account for the increased maxima seen in segments B and D. As a consequence we propose plage activity to increase the segment B and D maxima.

Flaring activity seems more prevalent in segment B. As seen in Figure 14, three of four flares are grouped near phase $\phi = 0.8$. Association with this deep minimum might imply some connection of flaring activity with the largest or coolest star spot, which is also at the same longitude as one of the two smaller spots seen best in segments A and C. The remaining flare, F4, lies close to $\phi = 0.2$, the other spot of the low-amplitude pair of spots. Spot/flare association was previously noted in the M dwarf EV Lac, shown also to have longitude-dependent flaring associated with a star spot site (Leto et al., 1997). However, for EV Lac, flares were detected a year before the spots became easily detectable by their system ($\Delta V \sim 0.1$) and, once spots formed, flare activity abated.

5.2. A Small, Variable Spot on Barnard’s Star

The periodogram (Figure 12) does not provide a clear identification of a single period of variation. The trend-corrected direct light curve (Figure 16, top) has been fit with a sin wave, constraining $P = 130^d.4$. The constant amplitude is $\Delta V \sim 0^m.01$, about five times our formal photometric error. Given the sparse coverage, it is speculative to interpret this light curve as showing rotational modulation of a single, small spot decreasing in size.

5.3. Rotation Periods for Proxima Cen and Barnard’s Star

Rotation periods for Proxima Cen have been predicted from chromospheric activity levels by Doyle (1987), who obtained $P = 51^d \pm 12^d$. Guinan & Morgan (1996) measure a rotation period ($P = 31^d.5 \pm 1^d.5$) from IUE observations of strong Mg II h+k emission at 280nm. We find no support for either rotation period in our periodogram (Figure 11) or light curves, direct (Figures 13) or phased (Figure 14). We do note that the variation due to the spot pair produces a period between the Doyle prediction and Guinan & Morgan measurement.

The observed variations for Proxima Cen and Barnard’s Star, if interpreted as rotationally modulated spots, yield rotation periods far longer than for other M stars. For example, Bouvier et al., 1995 find $4^d < P < 8^d$ for a sample of young, early-M T Tauri stars. Magnetic braking is postulated to slow rotation over time (Cameron & Campbell, 1993). The inferred rotation period for Proxima Cen is consistent with old age. This age may be 4 - 4.5 By, if Proxima Cen is coeval with α Cen (Demarque et al., 1986). A relatively older age for Barnard’s Star can be surmised from lower than solar metallicity (Gizis, 1996) and higher than solar space velocity (Eggen, 1996), both consistent with a longer rotation period, if one accepts the reality of the variation.

5.4. Shorter Time-scale Variations

The level of internal per-orbit precision for these photometric data is near 0.002 magnitudes. Hence, the dispersion about the phased light curves for Proxima Cen (Figure 14) is likely intrinsic to the stars. Two possibilities are miniflaring and the creation and destruction of small star spots and plages. That either phenomenon must have a duration longer than hours, at least for Proxima Cen, is suggested by the segment A phased light curve (Figure 14, bottom) and a detailed light curve for segment A (see Benedict et al., 1993, Fig. 3). Segment A contains four pairs of back-to-back orbits and one set of three contiguous orbits (on mJD 8845). In each case the time on target coverage is over 90 minutes. For most of these contiguous orbits differences are within two standard deviations and not statistically significant. Since ‘flare’ implies a relatively short duration, miniflaring cannot be the cause of the scatter.

5.5. Activity Cycles

The $\sim 1100^d$ cycle of alternating high and low amplitude (see Figure 13) is suggestive of an activity cycle for Proxima Cen. However, the gap in our coverage in segment C weakens any claims that can be made relative to the timing of this cycle. Comparing their 1995 IUE data with earlier archival data, Guinan & Morgan, 1996 propose an activity cycle that was in a low-state in 1995, agreeing with our identification of segment C representing a low-state (Figure 13).

6. Conclusions

1. For FGS 3 photometry we have identified four sources of systematic error: background contamination (primarily Zodiacal Light); spatial flat field variations (significant only for target positions $r > 20''$ from pickle center); temporal sensitivity changes (calibrated to a level introducing a 0.001 magnitude differential run-out error in 1000 days); and a possible warm-up effect (see section 3.3).

2. Two to four short ($t \leq 100^s$) observations with FGS 3 during one orbit yield 2 milli-magnitudes precision photometry, provided the targets are bright ($V \leq 11.0$) and restricted to the central $20''$ of the pickle.

3. Proxima Cen exhibits four distinct segments with two distinct behavior modes: short period, low amplitude and long period, large amplitude. These variations are consistent with a fundamental rotation period, $P = 83^d.5$ and three darker spots. Two of the spots are either very small or very low-contrast. They are spaced by 180° and persist throughout our temporal coverage, over 24 rotations. A single more prominent spot (either large or high-contrast) formed in less than one rotation period, persisted through four rotations, then disappeared. A spot reappeared within 5° of this same longitude five rotations later. The hemisphere opposite this

spot brightened each time the spot formed.

It is intriguing that active longitudes of spot formation separated by 180° are observed in chromospherically active stars with close stellar companions (Henry et al., 1995). If the photometric behavior of Proxima Cen is indicative of a synchronously rotating companion, its mass is less than that of Jupiter (Benedict et al., 1997a).

4. We interpret the four distinct segments with two distinct behavior modes seen in the Proxima Cen photometry as an activity cycle with a period $\sim 1100^d$. Most of the flare activity occurred in the long period, high amplitude variation segment B. In the phased lightcurve three of the four detected flares are near the deepest minimum.

5. The scatter in the Proxima Cen phased light curve is far larger than our photometric precision. This scatter could be caused by the formation and dissolution of small spots or plages within one rotation period.

6. We find brightness variations five times our formal photometric precision for Barnard’s Star. Unfortunately, the sparse coverage of the possible variation renders it a marginal detection. We conclude that Barnard’s Star shows very weak evidence for periodicity on a timescale of approximately 130 days

To confirm the spots and the inferred rotation periods will require observations of color changes (e.g. Vrba et al., 1988) and additional spectroscopic observations of lines sensitive to presence or absence of star spots. Extended-duration milli-magnitude V band photometry from the ground, while difficult (Gilliland et al., 1993), could probe the activity cycle periodicity of Proxima Cen. Future tests could include *Space Interferometry Mission* (Shao, 1995) observations with several microarcsec astrometric precision. If spots and plages exist on these stars, we can expect easily detectable star position shifts as activity sites vary. Such observations will provide detailed maps of spot and plage location. Extended temporal monitoring will provide evolutionary details.

Support for this work was provided by NASA through grants GTO NAG5-1603 and GO-06036.01-94A from the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. We thank Bill Spiesman and Artie Hatzes for discussions and draft paper reviews and Melody Brayton for paper preparation assistance. Denise Taylor provided crucial scheduling assistance at the Space Telescope Science Institute. Suggestions by an anonymous referee produced substantial improvements to the original draft.

Appendix 1

Appendix 1 contains the observation logs and measured average S values for Proxima Cen and Barnard’s Star.

REFERENCES

- Allen, C. W. 1983. "Astrophysical Quantities", The Athlone Press, London
- Bradley, A. J. et al. 1991. PASP, 103, 317
- Benedict, G. F. et al. 1993. PASP, 105, 487
- Benedict, G. F. et al. 1994a. PASP, 106, 327
- Benedict, G. F. et al. 1994b. BAAS, 185, 45.11
- Benedict, G. F. et al. 1997a. BAAS, 29, 1316
- Benedict, G. F. et al. 1997b. to appear in Proceedings of Cool Stars 10, Cambridge, MA
- Bouvier, J., Covino, E., Kovo, O. Martin, E. L., Matthews, J. M., Terranegra, L., and Beck, S. C. A&A, 299, 89
- Cameron, A. C. & Campbell, C. G. 1993. A&AS, 274, 309
- Cudworth, K., 1971. AJ, 76, 475
- Demarque, P., Guenther, D., & van Altena, W. 1986. ApJ, 300, 773
- Doyle, J. G. 1987. MNRAS, 224, 1p
- Eggen, O. J. 1996. AJ, 111, 466
- Gilliland, R. L. et al. 1993. AJ, 106, 2441
- Gilliland, R. L. & Dupree, A. K. 1996. ApJ, 463L, 29
- Gizis, J. E. 1997. AJ, 113, 806
- Gliese & Jahreiss 1991. Nearby Stars Catalog. NASA CDROM
- Guinan, E. F. & Morgan, N. D. 1996. BAAS, 188, 71.05
- Hatzes, A., 1993. ApJ, 410, 777
- Henry, G. W. , Eaton, J. A., Hamer, J. and Hall, D. S. 1995. ApJS, 97, 513
- Henry, T. J. and McCarthy, D. 1993. AJ, 106, 773
- Kirkpatrick, J. D. & McCarthy, D. 1994. AJ, 107, 333
- Kristian, J. 1991. Nature, 349, 747
- Kron, J. 1952. ApJ, 115, 301

- Liebert, J. & Probst, R. 1987. ARA&A, 25, 473
- Leto, G., Pagano, I., Buemi, C. S., & Rodono, M. 1997. A&A, 327, 1114
- Neff, J. E., O’Neal, D. & Sarr, S. H. 1995. ApJ, 452, 879
- Neff, J. E., Pagano, I., Rodono, M., Brown, A., Dempsey, R. C., Fox, D. C., & Linsky, J. L. 1996. A&A, 310, 173
- Panagi, P. M. & Mathioudakis, M. 1993. A&AS, 100, 343
- Pettersen, B., Hawley, S. L., & Fisher, G. 1992. Solar Physics, 142, 197
- Press et al., 1992. "Numerical Recipes in Fortran", 2nd edition, ch. 13.8
- Shao, M., 1995. BAAS, 187, 71.03
- Vrba, F. J., Herbst, W., & Booth, J. F. 1988. AJ, 96, 1032
- Walker, A. R. 1981. MNRAS, 195, 1029
- Whipple, A., McArthur, B. et al. 1995. Proc. *Hubble Space Telescope* Post-servicing Mission Calibration Workshop, A. Koratkar & C. Leitherer, eds. STScI. p. 119
- Zirin, H. "Astrophysics of the Sun", 1988. Cambridge University Press, Cambridge UK

Table 1. Proxima Cen

Parameter	Value	Reference
aliases	α Cen C, GJ 551, V645 Cen	
M_V	15.45 ± 0.1	
$B - V$	1.94	
Sp.T.	M5Ve	Gliese & Jahreiss, 1991
M_{Prox}	$0.11 M_{\odot}$	Kirkpatrick & McCarthy 1994
L_{Prox}	$0.001 L_{\odot}$	Liebert & Probst 1987
R_{Prox}	$0.15 R_{\odot}$	Panagi & Mathioudakis 1993

Table 2. Barnard’s Star

Parameter	Value	Reference
aliases	GJ 699, G 140-24, LHS 57	
M_V	13.2 ± 0.1	Gliese & Jahreiss, 1991
$B - V$	1.73	
Sp.T.	M4Ve	Kirkpatrick & McCarthy 1994
M_{Barn}	$0.17 M_{\odot}$	Henry & McCarthy, 1993
L_{Barn}	$0.0046 L_{\odot}$	Henry & McCarthy, 1993
R_{Barn}	$0.17 R_{\odot}$	

Table 3. Flat Field Modeling - M35 Stars

Parameter	S9_79	S11_68	S9_47
linear trend			
$I_o(counts)$	1.134	1.107	1.129
	± 0.008	0.023	0.015
Γ (counts d^{-1})	$-1.36E-05$	$-1.11E-05$	$-1.31E-05$
	0.10E-05	0.24E-05	0.10E-05
Variation from 180 deg flip			
A (counts)	0.0006	-0.0065	0.0039
	0.0006	0.0003	0.0004
P (days)	365.25	365.25	365.25
	0	0	0
ϕ (radians)	281.56	281.71	282.00
	1.03	0.17	0.14

Table 4. Flat Field Modeling - M Dwarf Stars

Parameter	GJ 623 and GJ 748
$I_o(counts)$	1.133 ± 0.0132
$I' (counts\ d^{-1})$	$-1.30E-05$ $0.13E-05$

Table 5. Proxima Cen Light Curve Parameters

Segment	N	I_o (counts)	I' (counts d ⁻¹)	A (counts)	P (days)	ϕ (radians)
A	32	3560	0.05	34.2	41.8	−1.6
		±710	0.08	6.0	0.9	2.7
BCD	39	3200	0.09	−102	82.5	−0.4
		160	0.02	12	0.3	2.4
B	17	3070	0.10	−139	82.7	−2.8
		580	0.06	9.5	0.7	0.7
C	11	4250	−0.02	−34.5	42.4	−2.9
		200	0.02	8.0	0.2	1.3
D	11	5473	−0.14	−114.8	84.3	−1.5
		645	0.06	16.3	0.8	2.2




Fig. 1.— The Barnard's Star field on 4 August 1994. North is at top, east to the left. The pickle coordinate system (x, y) is indicated. The symbol size is proportional to the relative brightness of each reference star (listed). The central circle (diameter $\sim 3''.8$) is accessible by the FGS 3 instantaneous aperture for any HST roll. Barnard's Star is labeled at three epochs.




Fig. 2.— S (average counts per 0.025 sec) for two faint astrometric reference stars in the Barnard’s Star field (stars 32 and 33, Figure 1) vs. target - Sun separation in degrees. Fitting function is equation 1.




Fig. 3.— Correcting for Zodiacal Light. Average intensity (S , counts per 0.025 sec) vs. modified Julian Date for the average of two faint astrometric reference stars in the Barnard’s Star field (Figure 2). The horizontal line denotes the mean brightness of the corrected data. Scatter is far less in the corrected data. Uncorrected and corrected values have been flat-fielded with equation 2




Fig. 4.— Field of view of FGS 3 with position of three M35 stars. The V magnitude is given as part of the identification (e.g., S9_47 has $V = 9.47$).




Fig. 5.— Time-dependent photometric variations for three stars in M35 modeled as linear trends whose parameters (I and I') are given in Table 3. The residuals in the top panels are size-coded to show the two LTSTAB orientations (Fall = small, Spring = large) and fit with sine functions. Note differences in the variation amplitude (parameter A in Table 3).




Fig. 6.— Trend of normalized intensity for Gl 623 and GJ 748. The fitted line parameters are given in Table 3




Fig. 7.— Amplitude of side-to-side variation vs. distance from pickle center. Targets are stars from M35 (location from Figure 4 and amplitude from Figure 5), stars from the Barnard’s Star reference frame (Figure 1), and GJ 748. Error bars along the X-axis indicate the radial range within the pickle for all observations of the particular target.




Fig. 8.— Intensities of Barnard field reference stars 34 and 36 plotted against pickle coordinates for all observation dates. The symbol size is proportional to the normalized intensity. The low-sensitivity region is clearly seen in the vicinity of $(x, y) = (-25, 627)$.




Fig. 9.— Photometry of Proxima Cen. Each orbit contains 2, 3, or 4 exposures (Appendix 1.1). Exposures perturbed by flares are marked F1 - F4. Exposures acquired during CVZ orbits are labeled 'cz'. Error bars are about the size of the plotted symbols. Four segments and two distinct behavior modes are identified, A - D. Trend line is discussed in text.




Fig. 10.— Photometry of Barnard’s Star. Each symbol represents the average of an exposure. Each orbit contains 3 exposures (Appendix 1.2). No flares were detected. Error bars are smaller than the plotted symbols. The time scale is identical to Figure 9. Trend line is discussed in text.




Fig. 11.— Periodogram from 71 normal points (average for each orbit) with flares removed, flat-fielded with equation 2. The periodogram has most significant peak at $P \sim 83$ days, with less than a 0.1% false-positive probability.




Fig. 12.— Periodogram from 35 normal points (average for each orbit), flat-fielded with equation 2. Periodogram has most significant peak at $P = 130.4^d$, with a 10% false-positive probability.




Fig. 13.— Direct light curve. Each symbol represents the average of from two to four exposures per orbit, with flares removed. Error bars are $1 - \sigma$. See Table 5, lines 1, 3, 4, and 5 for the results of fitting a sin wave and trend (equation 4) to each segment.




Fig. 14.— Phased light curves for Proxima Cen. Top: the two long-period segments (B and D, Figure 13). Sin wave fit has a period constrained to one cycle. Bottom: short-period segments (A and C) are phased to the longer period and show a double sin wave. The fit is constrained to have a period of one-half cycle. Error bars are about the size of the symbols. The observed flares are labeled F1 - F4.

Proxima Cen Phased Light Curve Low-amplitude, Short-period Signature in High-amplitude, Long-period Variations




Fig. 15.— Phased light curve for Proxima Cen. Bottom: the two long-period segments (B and D, Figure 13), phased to $P = 83.5^{\text{d}}$ and fit with a sin wave having a period of one cycle. Top: the residuals to the sin wave fit. These show a double sin wave pattern nearly identical to the two short-period segments, A and C (Figures 13 and 14), suggesting that the low-amplitude, short-period signature persists during segments B and D.




Fig. 16.— Per-orbit average direct light curve. Each symbol represents the average of three exposures. Error bars are $1 - \sigma$. Bottom: fit with a trend line. Top: residuals to trend line, fit with a constant-amplitude sin wave with constrained $P = 130^{\text{d}}.4$.




Fig. 17.— Trend-corrected light curve for Barnard’s Star, phased to $P = 130.4^d$. The full-amplitude of the variation is ~ 0.01 magnitude. Error bars are $1 - \sigma$.